

## SHRIMP U-Pb geochronology of the Choma-Kalomo block (Zambia) and geological implications

Bulambo M. <sup>1</sup>, De Waele B. <sup>2</sup>, Kampunzu A.B. <sup>3</sup>, Tembo F. <sup>4</sup>

<sup>1</sup> Department of Geology, University of Lubumbashi, D.R.Congo (jbulambo@yahoo.fr)

<sup>2</sup> School of Applied Geology, Curtin University of Technology, WA 6102 Australia

<sup>3</sup> Department of Geology, University of Botswana, Private Bag 0022, Gaborone, Botswana

<sup>4</sup> Department of Geology, School of Mines, University of Zambia, PO Box 32 379, Lusaka, Zambia

The Choma-Kalomo block is a Mesoproterozoic terrane exposed in southern Zambia, south of the Pan-African Zambezi belt. It includes a large plutonic complex in the central part, surrounded by supracrustal metasedimentary assemblages. The plutonic complex represents a set of intrusive bodies that yielded TIMS U-Pb zircon dates between c. 1345 and 1200 Ma (Hanson et al., 1988). Two deformation events were recorded in this terrane: D1 is presumably coeval with the emplacement of the c.1345 Ma granitoids (converted into orthogneisses) and D2 coeval with emplacement of c.1200 Ma granitoids.

Because of its location between the Mesoproterozoic Irumide belt of Zambia and the Ngamiland province of Botswana, the Choma-Kalomo block is a critical terrane in the reconstruction of late-Mesoproterozoic (Kibaran) orogenic system central and southern Africa.

Zircons extracted from five granitoid samples were investigated by U-Pb SHRIMP technique and yielded the following results: CK4: medium-grained two mica granite: 1174±27 Ma; CK10: foliated medium to coarse biotite granite: 1188±11Ma; CK12: gneiss containing mica-rich restites: 1177±70 Ma; CK25: foliated, medium to coarse grained biotite granite: 1181±9 Ma; CK13: augen (ortho)gneiss: 1368±10 Ma.

The new SHRIMP ages support the existence of two major magmatic events in the Choma-Kalomo block at c. 1.37 and 1.18 Ga. Neither the 1.37 Ga magmatic event nor the 1.18 event are reported from the Irumide belt (De Waele et al., 2002) or the Ngamiland province (Kampunzu et al., 1999), which is not in support of the correlation of the Choma-Kalomo block to these two provinces as suggested by Hanson et al. (1988) and Singletary et al. (2002). In contrast, 1.38-1.37 Ga granitoids are widespread in the Kibaran belt in SE and eastern Congo (Kokonyangi et al., 2001) and in Rwanda and Burundi (Wingate and De Waele, pers. comm.). A second (<1.25 and >1.0 Ga) and third (ca. 1 Ga tin granite) magmatic events described in the Kibaride belt include granitoids emplaced between <1.25-1.0 Ga (Kampunzu and Kokonyangi, pers. comm.). Therefore, we suggest that the Choma-Kalomo terrane is a block detached from the Kibaran belt of southeastern Congo during the rift-drift stage of the Katangan-Zambezi basin evolution.

If this interpretation is correct, it implies that either the Katangan-Zambezi basin did not evolve into an Atlantic-type Mid-Oceanic Rift (e.g. Kampunzu et al., 1991, 2000) or that the closure of the Neoproterozoic Katangan Ocean fortuitously reunited the dispersed blocks in their original, pre-drift position which is unlikely. The existence of HP metamorphic assemblages (e.g. John et al., 2003) along the Katangan-Zambezi belts demand substantial oceanic subduction and would argue for an extensive Katangan-Zambezi basin but how extensive was this basin is still unconstrained. The absence of Neoproterozoic arc igneous rocks in the Zambezi-Lufilian belts and the geochronological data in this paper indicating that the Choma-Kalomo block on the southern side of these two Neoproterozoic belts is a terrane detached from the Kibaride on the northern side of the same belt do not support the Neoproterozoic closure of an ATL-type oceanic basin in this region. These conflicting data require further studies before a definitive conclusion is reached.

### References

- De Waele, B., Mapani, B., 2002. Geology, stratigraphy and correlation of the Central Irumide belt. *JAES* 35, 385-397.
- Hanson, R.E., Wilson, T.J., Brueckner, H.K., Onstott, T.C., Wardlaw, M.S., Johns, C.C. and Hardcastle, K.C., 1988. Reconnaissance geochronology, tectonothermal evolution, and regional significance of the middle Proterozoic Choma-Kalomo block, southern Zambia. *Precambrian Research* 42, 39-61
- John, T., Schenk, V., Haase, K., Scherer, E., Tembo, F., 2003. Evidence for a Neoproterozoic ocean in south-central Africa from mid-oceanic-ridge-type geochemical signatures and pressure-temperature estimates of Zambian eclogites. *Geology* 31, 243-246.
- Kampunzu, A.B., Mohr, P., 1991. Magmatic evolution and petrogenesis in the East-African rift system. In: Kampunzu, A.B., Lubala, R.T. (eds) *Magmatism in extensional tectonic settings. The Phanerozoic African Plate*. Springer, Berlin Heidelberg New York, pp 85-136.
- Kampunzu, A.B., Tembo, F., Matheis, G., Kapenda, D., Huntsman-Mapila, P., 2000. Geochemistry and tectonic setting of mafic igneous units in the Neoproterozoic Katangan basin, central Africa: implications for Rodinia break up. *Gondwana Research* 3, 125-153.
- Kampunzu, A.B., Armstrong, R., Modisi, M.P., Mapeo, R.B., 1999. The Kibaran belt in southwest Africa: Ion microprobe U-Pb zircon data and definition of the Kibaran Ngamiland belt in Botswana, Namibia and Angola. *Journal of African Earth Sciences*, GSA 11 Special Abstract Issue 28, pp. 34..
- Kokonyangi, J., Okudaira, T., Kampunzu, A.B., Yoshida, M., 2001. Geological evolution of the Kibaride belt, Mitwaba, Democratic Republic of Congo, Central Africa. *Gondwana Research* 4, 663-664.
- Singletary, S.J., Hanson, R.E., Martin, M.W., Martin, M.W., Crowley, J.L., Bowring, S.A., Key, R.M., Ramokate, L.V., Direng, B.B., Krol, M.A., 2002. Geochronology of basement rocks in the Kalahari Desert, Botswana, and implication for regional Proterozoic tectonics. *Precambrian Research* 121, 47-71.

**Keywords:** Geochronology; Choma-Kalomo block; Mesoproterozoic; Kibaran belt.